



# LABORATOIRES EUROSILICONE

Siège social : Z.I. de la Peyrolière - B.P. 68 - 84402 APT - FRANCE  
Tél. : + 33 (0) 4 90 04 30 30 - Fax : + 33 (0) 4 90 04 60 06  
Email : contact@eurossilicone.com - Web : http://www.eurossilicone.com



## LIFETIME GUARANTEE

\* \* \*

All EUROSILICONE mammary prostheses are manufactured in accordance with ISO 9001/EN 46001 and CE mark. These certificates guarantee the quality and control in the production of Eurosilicone products.

EUROSILICONE Laboratories use only the highest quality medical grade silicone in the manufacture of mammary implants, supplied by the best FDA approved American supplier.

EUROSILICONE Laboratories are confident of their silicone gel filled mammary prostheses and have therefore decided to give a lifetime guarantee to the implanted gel filled mammary prostheses.

## LIFETIME GUARANTEE

EUROSILICONE Laboratories guarantee the free replacement of their silicone gel filled mammary prosthesis which have ruptured inside.

This guarantee applies to all mammary prosthesis filled with silicone gel, implanted after June 14<sup>th</sup> 98 by surgeons holding a recognised and appropriate qualification and carrying out implantation with respect to the hygiene norms as far as plastic mammary surgery is concerned. In case of any rupture the free replacement will be identical to that implant removed (reference & volume).

## REQUESTED CONDITIONS FOR THE IMPLEMENTATION OF LIFETIME GUARANTEE

The free lifetime guarantee, after a rupture of a silicone gel filled prosthesis will be valid only under the following conditions :

- Full respect of the traceability
- Return the explanted prosthesis with the attached EUROSILICONE document mentioning the reasons for the ex-plantation and the observations during the removal. It should be completed by the surgeon and co-signed by the patient

This guarantee covers all manufacturing defects leading to rupture of the prosthesis.

*All ruptures from outside causes are excluded from the guarantee.*

For instance, (this list not being exhaustive), all ruptures coming from : capsular contraction, bad manipulation of a surgical instrument during the implantation or surgery following trauma, or assault or any other injury to the patient.

\* \* \*